Trans 4 zlom 16.1.2008 12:15 Str. 215
proLékaře.cz | 24.11.2026

lymphoma. J Nucl Med 2000; 41: 952–958.
6. **Gordon LI, Witzig TE, Wiseman GA, et al.** Yttrium 90 ibritumomab tiuxetan radioimmunotherapy for relapsed or refractory low-grade non-Hodgkin's lymhoma. Sem Oncol 2002; 29 (Suppl 2): 87–92.
7. **Witzig TE, White CA, Wiseman GA, Gordon LI, Emmanouilides C, Raubitschek A, et al.** Phase I/II trial of IDEC-Y2B8 radioimmunotherapy for treatment of relapsed or refractory CD20+ B-cell non-Hodgkin's lymphoma. J Clin Oncol 1999; 17: 3793–3803.
8. **Witzig TE, Gordon LI, Cabanillas F, Czuczman MS, Emmanouilides C, Joyce R, et al.** Randomized controlled trial of yttrium-90-labeled ibritumomab tiuxetan radioimmunotherapy versus rituximab immunotherapy for patients with relapsed or refractory low-grade, follicular, or transformed B-cell non-Hodgkin's lymphoma. J Clin Oncol 2002; 20: 2453–2463.
9. **Gordon LI, Molina A, Witzig T, Emmanouilides C, Raubtischek A, Darif M, et al.** Durable responses after ibritumomab tiuxetan radioimmunotherapy for CD20+ B-cell lymphoma: long-term follow-up of a phase 1/2 study. Blood 2004; 103: 4429–4431.
10. **Ansell SM, Ristow KM, Hanermann TM, et al.** Subsequent chemotherapy regimens are well tolerated after radioimmunotherapy with yttrium-90 ibrituomomab tiuxetan fo non-Hodgkin's lymphoma. J Clin Oncol 2002; 20: 3885–3890.
11. **Emmanoulides C, Witzig TE, Gordon LI, et al.** Treatment with yttrium 90 ibritumomab tiuxetan at early relapse is safe and effective in patients with previously treated B-cell non-Hodgkin's lymphoma. Leuk Lymphoma 2006; 47: 629–636.
12. **Jacobs SA , Vidnovic N, Joyce J, et al.** Full-dose 90Y ibritumomab tiuxetan therapy is safe in patients with prior myeloablative chemotherapy. Clin Cancer Res 2005; 11: 7146–7150.
13. **Shipp MA, Harrington DP, Anderson JR, et al.** A predictive model for aggressive non-Hodgkin's lymphoma. N Engl J Med 1993; 329: 987–994.
14. **Solal-Céligny P, Roy P, Colombat P, et al.** Follicular lymphoma prognostic index. Blood 2004; 104: 1258–1265.
15. Cheson BD, Horning SJ, Coiffier B, et al. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomnas. J Clin Oncol 1999; 17: 1244–1253.
16. **Shipley DL, Spigel DR, Carell DL, et al.** Phase II trial of rituximab and short duration chemotherapy followed by 90Y ibritumomab tiuxetan as first-line treatment for patients with follicular lymphoma: a Minnie Pearl Cancer Research Network phase II trial. J Clin Oncol ASCO Annual Meeting Proceedings 2004; 22 (Suppl 14): 6519.
17. **Jurczak WJ, Szostek MS, Giza AG, et al.** Consolidation of chemotherapy response in mantle cell lymphoma (MCL) patients with 90Y-ibritumomab tiuxetan (90Y-Zevalin®) radioimmunotherapy (RIT). Haematologica 2006: 91 (Suppl. 1): 188.
18. **Dreyling M, Trümper L, von Schilling C, et al.** Results of national consensus workshop: therapeutic algorithm in patients with follicular lymphoma – role of radioimmunotherapy. Ann Hematol 2007; 86: 81–87.
19. **Rose AL, Smith B, Maloney DG.** Glucocorticoids and rituximab in vitro: synergistic direct antiproliferative and apoptotic effects. Blood 2002; 100: 1765– 1773.

Podpořeno grantem IGA MZ ČR No. NR/9502-3 a VZ MŠM 619 895 9205

MUDr. Tomáš Papajík, CSc.
Hemato-onkologická klinika LF UP a FN Olomouc
I. P. Pavlova 6
775 20 Olomouc
e-mail: tomas.papajik@fnol.cz

GRATULUJEME

Ve druhé polovině roku 2007 někteří členové Společnosti pro transfuzní lékařství ČLS JEP oslavili významná životní výročí. Všem kolegyním a kolegům gratulujeme a přejeme jim další cestu životem, naplněnou zdravím, štěstím a životní pohodou.

MUDr. Banzetová Helena
MUDr. Černohorská Jitka
MUDr. Fenclová Eva
prof. MUDr. Indrák Karel, DrSc.
Knopová Jitka
MUDr. Jitka Mačalová
doc. MUDr. Neuwirthová Radana, CSc.
Tlapáková Eva
MUDr. Tomancová Marie
MUDr. Valentová Kamila
Alena Vašková
MUDr. Zvarová Marta

Redakční rada časopisu Transfuze a hematologie dnes, výbor STL